

Alabama requires all students in grades 6-12 to have a Tdap vaccine.

In addition, ADPH and ACIP recommends HPV and Meningococcal vaccine for this age group.

For questions, please contact the Immunization Division at 1-800-469-4599.



ADPH.ORG



## **Important Vaccines for Your Adolescent**

The Alabama Department of Public Health (ADPH), Immunization Division (IMM), and the Advisory Committee of Immunization Practices (ACIP) encourage you to make sure that your child receives all recommended vaccines in order to prevent diseases. Read about these vaccines below.

## Immunizations recommended for persons 11 through 18 years of age<sup>a</sup>

Vaccination	Required or Recommended	For whom does the ACIP/CDC <sup>b</sup> recommend this vaccine?	What does this vaccine help protect against?	What are some of the symptoms of these diseases?	When is the vaccine given?
Tdap <sup>c</sup>	Required by State Law	• All adolescents 11 through 18 years of age	3 serious bacterial diseases     Tetanus: also known as "lockjaw"     Diphtheria: upper respiratory disease     Pertussis: also known as "whooping cough"	Tetanus: Painful tightening of the muscles  Diphtheria: Difficulty in breathing; can lead to paralysis, heart failure  Pertussis: Severe coughing spells and vomiting	• 1 dose during adolescence
Varicella	Required by State Law	• All children 12 months through 18 years of age	Chickenpox	A skin rash of blister-like lesions, usually on the face, scalp, or trunk	One dose required <sup>e</sup> Two doses recommended <sup>e</sup>
Meningococcal	Recommended	• All adolescents 11 through 18 years of age	Meningococcal disease including blood infection (sepsis) and meningitis	Initial symptoms can be flu-like (including fever), stiff neck, rash     Can lead to hearing loss, amputation, seizures, coma, shock, or death within 48 hours	1 dose at 11 through 12 years of age     1 booster dose at 16 years of age     or     1 dose at 13 through 18 years of age if not previously vaccinated     1 booster dose at age 16 through 18 years for persons who received 1st dose at 13 through 15 years of age
HPV <sup>d</sup>	Recommended	• All females and males 11 through 26 years of age	Sexually transmitted infection caused by the human papillomavirus     Can cause cervical cancer or genital warts	• No symptoms at first	
Influenza (Seasonal)	Recommended	• All children 6 months through 18 years of age	• Influenza virus: the "flu"	• Fever, sore throat, cough, head and muscle aches, extreme tiredness	One dose each year Children 6 months—8 years of age: 2 doses may be needed 4 weeks apart

Other vaccines recommended include Hepatitis A and Hepatitis B.

The above vaccination information and additional immunization facts are available from the CDC online at: <a href="http://www.cdc.gov/vaccines/spec-grps/preteens-adol.htm">http://www.cdc.gov/vaccines/spec-grps/preteens-adol.htm</a> or <a href="http://www.adph.org/immunization/">http://www.adph.org/immunization/</a> or contact ADPH Immunization at 1-800-469-4599.

<sup>&</sup>lt;sup>a</sup> Additional vaccines may be recommended depending on the child's vaccination history and risk factors. Please see the complete vaccination schedule at: <a href="http://www.cdc.gov/vaccines/recs/schedules/child-schedule.htm">http://www.cdc.gov/vaccines/recs/schedules/child-schedule.htm</a>

<sup>&</sup>lt;sup>b</sup> CDC = Centers for Disease Control and Prevention.

<sup>&</sup>lt;sup>c</sup> Tdap = Tetanus, diphtheria, and acellular pertussis.

<sup>&</sup>lt;sup>d</sup> HPV = Human papillomavirus vaccine. ACIP recommends that all males and females ages 11 through 12 years be routinely vaccinated with a 3-dose series.

<sup>&</sup>quot;See the School Immunization Law at www.adph.org/immunization